Brain Research, 553 (1991) 358–360
Elsevier

Author Index

Acciarri, N., see Morari, M., 14
Adams, F., Schwarting, R.K.W., Boix, F. and Huston, J.P., Lateralized changes in behavior and striatal dopamine release following unilateral tactile stimulation of the perioral region: a microdialysis study, 318
Altar, C.A., see Baudry, M., 215
Amiri', Z., Weizman, R., Katz, Y., Burstein, O., Edoute, Y., Lochner, A. and Gavish, M., Testosterone and cyproterone acetate modulate peripheral but not central benzodiazepine receptors in rats, 155
Antonelli, T., see Morari, M., 14
Araki, T., see Kato, H., 238
Araki, T., see Kato, H., 33
Asikainen, A., see Ylinen, A., 195

Baird, A., see Frautschy, S.A., 291
Barbaresi, P., see Minciacchi, D., 309
Baudry, M. and Altar, C.A., Entorhinal cortex lesion or intrahippocampal colchicine injection increases peripheral type benzodiazepine binding sites in rat hippocampus, 215
Beani, L., see Morari, M., 14
Becker, C.G., see Becker, T., 331
Becker, T. and Becker, C.G., Cell migration along glial fibers in dissociated cell culture of the frog optic tectum, 331
Berthoud, H.-R. and Powley, T.L., Morphology and distribution of efferent vagal innervation of rat pancreas as revealed with anterograde transport of DiI, 336
Besson, J.M., see Puig, S., 222
Bhargava, H.N., see Gudehithlu, K.P., 284
Bianchi, C., see Morari, M., 14
Bignami, A., see Dahl, D., 163
Bishai, I., see Hynes, N., 4
Black, I.B., see Pincus, D.W., 211
Boix, F., see Adams, F., 318
Bondoux, D., see Chapouthier, G., 342
Braun, A., see Gibson, R.E., 110
Brooks, P., see Lee, K.S., 171
Burstein, O., see Amiri', Z., 155

Calo, G., see Morari, M., 14
Carey, F., see Strijbos, P.J.L.M., 249
Carman, L.S., Gage, F.H. and Shults, C.W., Partial lesion of the substantia nigra: relation between extent of lesion and rotational behavior, 275
Chalmers, D.T., see Dewar, D., 58
Chapouthier, G., Bondoux, D., Martin, B., Desforges, C. and Launay, J.-M., Genetic difference in sensitivity to β-carboline: evidence for the involvement of brain benzodiazepine receptors, 342
Coceani, F., see Hynes, N., 4
Corp, E.S. and Smith, G.P., Characterization of axonally transported [^{125}I]-PYY binding sites in rat vagus nerve, 175
Cory-Slechta, D.A. and Widzowski, D.V., Low level lead exposure increases sensitivity to the stimulus properties of dopamine D_1 and D_2 agonists, 65

Dahl, D. and Bignami, A., Neurofilament phosphorylation in the axonless horizontal cells of rat retina, 163
Dauphin, F., Lacombe, P., Sercombe, R., Hamel, E. and Seylaz, J., Hypercapnia and stimulation of the substantia innominata increase rat frontal cortical blood flow by different cholinergic mechanisms, 75
Davar, G., Hama, A., Deykin, A., Vos, B. and Maciewicz, R., MK-801 blocks the development of thermal hyperalgesia in a rat model of experimental painful neuropathy, 327
Demura, H., see Nakano, Y., 1
Desforges, C., see Chapouthier, G., 342
Dewar, D., Chalmers, D.T., Graham, D.I. and McCulloch, J., Glutamate metabotropic and AMPA binding sites are reduced in Alzheimer's disease: an autoradiographic study of the hippocampus, 58
Deykin, A., see Davar, G., 327
Diciccobloom, E., see Pincus, D.W., 211

Edoute, Y., see Amiri', Z., 155

Fabrizi, A., see Morari, M., 14
Fattaccini, C.-M., see Orer, H.S., 123
Forder, R., see Strijbos, P.J.L.M., 249
Frautschy, S.A., Walicke, P.A. and Baird, A., Localization of basic fibroblast growth factor and its mRNA after CNS injury, 291
Frenk, H., see Schul, R., 353
Freund, T.F., see Ylinen, A., 195
Fujimori, K., see Nakazawa, K., 44

Gackenheimer, S.L., see Gehlert, D.R., 149
Gage, F.H., see Carman, L.S., 275
Gaist, G., see Morari, M., 14
Gavish, M., see Amiri', Z., 155
Gehlert, D.R., Gackenheimer, S.L., Wong, D.T. and Robertson, D.W., Localization of 5-HT$_3$ receptors in the rat brain using [^{3}H]LY278584, 149
Gibson, R.E., Zeeberg, B.R., Melograna, J.M., Wang, T.F., Ruch, J., Braun, A. and Reba, R.C., In vivo dissociation kinetics of [^{3}H]quinuclidinyl benzilate: relationship to muscarinic receptor concentration and in vitro kinetics, 110
Gonzalez-Mora, J.L., see Louilot, A., 313
Graham, D.I., see Dewar, D., 58
Granato, A., see Minciacchi, D., 309
Guadalupe, T., see Louilot, A., 313
Gudehithlu, K.P., Tejwani, G.A. and Bhargava, H.N., β-Endorphin and methionine-enkephalin levels in discrete brain regions, spinal cord, pituitary gland and plasma of morphine tolerant–dependent and abstinent rats, 284
Gulyas, A., see Ylinen, A., 195

Hall, E.D. and Smith, S.L., The 21-aminosteroid antioxidant tirilazad mesylate, U-74006F, blocks cortical hypoperfusion following spreading depression, 243
Hama, A., see Davar, G., 327
Hamel, E., see Dauphin, F., 75
Hara, H., see Kato, H., 33
Hexum, T.D., see Li, W., 167
Hsiao, C.-F., see Huang, C., 129
Huang, C., Liu, G., Hsiao, C.-F. and Huang, R.H., Nitrous oxide inhibits auditory and visual responses of granule cells in the cerebellum, 129
Huang, C.-m. and Liu, G., Auditory responses in the posterior vermis of the cat: the buried cerebellar cortex, 201
Huang, R.H., see Huang, C., 129
Huston, J.P., see Adams, F., 318
Hynes, N., Bishai, I., Lees, J. and Coceani, F., Leukotrienes in brain: natural occurrence and induced changes, 4

Ihara, N., Ueda, S., Tanabe, T. and Sano, Y., Transplantation of fetal mesencephalic and medullary raphe tissues into the cerebellum of denervated adult rats – an immunohistochemical study, 39
Imagawa, M., see Sasaki, S.-I., 159
Inoue, K., see Nakazawa, K., 44

Jope, R.S., see Ormandy, G.C., 51

Kalasapudi, V.D., see Weiner, E.D., 117
Kaneko, S., see Matsuoka, N., 188
Kangrga, I. and Randić, M., Outflow of endogenous aspartate and glutamate from the rat spinal dorsal horn in vitro by activation of low- and high-threshold primary afferent fibers. Modulation by μ-opioids, 347
Kato, H., Araki, T., Hara, H. and Kogure, K., Sequential changes in muscarinic acetylcholine, adenosine A_1 and calcium antagonist binding sites in the gerbil hippocampus following repeated brief ischemia, 33
Kato, H., Liu, Y., Araki, T. and Kogure, K., Temporal profile of the effects of pretreatment with brief cerebral ischemia on the neuronal damage following secondary ischemic insult in the gerbil: cumulative damage and protective effects, 238
Katz, Y., see Amiri', Z., 155
Kincaid, A.E., Penney Jr., J.B., Young, A.B. and Newman, S.W., The globus

Brain Research, 553 (1991) 358–360
Elsevier

Author Index

Acciarri, N., see Morari, M., 14
Adams, F., Schwarting, R.K.W., Boix, F. and Huston, J.P., Lateralized changes in behavior and striatal dopamine release following unilateral tactile stimulation of the perioral region: a microdialysis study, 318
Altar, C.A., see Baudry, M., 215
Amiri', Z., Weizman, R., Katz, Y., Burstein, O., Edoute, Y., Lochner, A. and Gavish, M., Testosterone and cyproterone acetate modulate peripheral but not central benzodiazepine receptors in rats, 155
Antonelli, T., see Morari, M., 14
Araki, T., see Kato, H., 238
Araki, T., see Kato, H., 33
Asikainen, A., see Ylinen, A., 195

Baird, A., see Frautschy, S.A., 291
Barbaresi, P., see Minciacchi, D., 309
Baudry, M. and Altar, C.A., Entorhinal cortex lesion or intrahippocampal colchicine injection increases peripheral type benzodiazepine binding sites in rat hippocampus, 215
Beani, L., see Morari, M., 14
Becker, C.G., see Becker, T., 331
Becker, T. and Becker, C.G., Cell migration along glial fibers in dissociated cell culture of the frog optic tectum, 331
Berthoud, H.-R. and Powley, T.L., Morphology and distribution of efferent vagal innervation of rat pancreas as revealed with anterograde transport of DiI, 336
Besson, J.M., see Puig, S., 222
Bhargava, H.N., see Gudehithlu, K.P., 284
Bianchi, C., see Morari, M., 14
Bignami, A., see Dahl, D., 163
Bishai, I., see Hynes, N., 4
Black, I.B., see Pincus, D.W., 211
Boix, F., see Adams, F., 318
Bondoux, D., see Chapouthier, G., 342
Braun, A., see Gibson, R.E., 110
Brooks, P., see Lee, K.S., 171
Burstein, O., see Amiri', Z., 155

Calo, G., see Morari, M., 14
Carey, F., see Strijbos, P.J.L.M., 249
Carman, L.S., Gage, F.H. and Shults, C.W., Partial lesion of the substantia nigra: relation between extent of lesion and rotational behavior, 275
Chalmers, D.T., see Dewar, D., 58
Chapouthier, G., Bondoux, D., Martin, B., Desforges, C. and Launay, J.-M., Genetic difference in sensitivity to β-carboline: evidence for the involvement of brain benzodiazepine receptors, 342
Coceani, F., see Hynes, N., 4
Corp, E.S. and Smith, G.P., Characterization of axonally transported [^{125}I]-PYY binding sites in rat vagus nerve, 175
Cory-Slechta, D.A. and Widzowski, D.V., Low level lead exposure increases sensitivity to the stimulus properties of dopamine D_1 and D_2 agonists, 65

Dahl, D. and Bignami, A., Neurofilament phosphorylation in the axonless horizontal cells of rat retina, 163
Dauphin, F., Lacombe, P., Sercombe, R., Hamel, E. and Seylaz, J., Hypercapnia and stimulation of the substantia innominata increase rat frontal cortical blood flow by different cholinergic mechanisms, 75
Davar, G., Hama, A., Deykin, A., Vos, B. and Maciewicz, R., MK-801 blocks the development of thermal hyperalgesia in a rat model of experimental painful neuropathy, 327
Demura, H., see Nakano, Y., 1
Desforges, C., see Chapouthier, G., 342
Dewar, D., Chalmers, D.T., Graham, D.I. and McCulloch, J., Glutamate metabotropic and AMPA binding sites are reduced in Alzheimer's disease: an autoradiographic study of the hippocampus, 58
Deykin, A., see Davar, G., 327
Diciccobloom, E., see Pincus, D.W., 211

Edoute, Y., see Amiri', Z., 155

Fabrizi, A., see Morari, M., 14
Fattaccini, C.-M., see Orer, H.S., 123
Forder, R., see Strijbos, P.J.L.M., 249
Frautschy, S.A., Walicke, P.A. and Baird, A., Localization of basic fibroblast growth factor and its mRNA after CNS injury, 291
Frenk, H., see Schul, R., 353
Freund, T.F., see Ylinen, A., 195
Fujimori, K., see Nakazawa, K., 44

Gackenheimer, S.L., see Gehlert, D.R., 149
Gage, F.H., see Carman, L.S., 275
Gaist, G., see Morari, M., 14
Gavish, M., see Amiri', Z., 155
Gehlert, D.R., Gackenheimer, S.L., Wong, D.T. and Robertson, D.W., Localization of 5-HT$_3$ receptors in the rat brain using [^{3}H]LY278584, 149
Gibson, R.E., Zeeberg, B.R., Melograna, J.M., Wang, T.F., Ruch, J., Braun, A. and Reba, R.C., In vivo dissociation kinetics of [^{3}H]quinuclidinyl benzilate: relationship to muscarinic receptor concentration and in vitro kinetics, 110
Gonzalez-Mora, J.L., see Louilot, A., 313
Graham, D.I., see Dewar, D., 58
Granato, A., see Minciacchi, D., 309
Guadalupe, T., see Louilot, A., 313
Gudehithlu, K.P., Tejwani, G.A. and Bhargava, H.N., β-Endorphin and methionine-enkephalin levels in discrete brain regions, spinal cord, pituitary gland and plasma of morphine tolerant–dependent and abstinent rats, 284
Gulyas, A., see Ylinen, A., 195

Hall, E.D. and Smith, S.L., The 21-aminosteroid antioxidant tirilazad mesylate, U-74006F, blocks cortical hypoperfusion following spreading depression, 243
Hama, A., see Davar, G., 327
Hamel, E., see Dauphin, F., 75
Hara, H., see Kato, H., 33
Hexum, T.D., see Li, W., 167
Hsiao, C.-F., see Huang, C., 129
Huang, C., Liu, G., Hsiao, C.-F. and Huang, R.H., Nitrous oxide inhibits auditory and visual responses of granule cells in the cerebellum, 129
Huang, C.-m. and Liu, G., Auditory responses in the posterior vermis of the cat: the buried cerebellar cortex, 201
Huang, R.H., see Huang, C., 129
Huston, J.P., see Adams, F., 318
Hynes, N., Bishai, I., Lees, J. and Coceani, F., Leukotrienes in brain: natural occurrence and induced changes, 4

Ihara, N., Ueda, S., Tanabe, T. and Sano, Y., Transplantation of fetal mesencephalic and medullary raphe tissues into the cerebellum of denervated adult rats – an immunohistochemical study, 39
Imagawa, M., see Sasaki, S.-I., 159
Inoue, K., see Nakazawa, K., 44

Jope, R.S., see Ormandy, G.C., 51

Kalasapudi, V.D., see Weiner, E.D., 117
Kaneko, S., see Matsuoka, N., 188
Kangrga, I. and Randić, M., Outflow of endogenous aspartate and glutamate from the rat spinal dorsal horn in vitro by activation of low- and high-threshold primary afferent fibers. Modulation by μ-opioids, 347
Kato, H., Araki, T., Hara, H. and Kogure, K., Sequential changes in muscarinic acetylcholine, adenosine A_1 and calcium antagonist binding sites in the gerbil hippocampus following repeated brief ischemia, 33
Kato, H., Liu, Y., Araki, T. and Kogure, K., Temporal profile of the effects of pretreatment with brief cerebral ischemia on the neuronal damage following secondary ischemic insult in the gerbil: cumulative damage and protective effects, 238
Katz, Y., see Amiri', Z., 155
Kincaid, A.E., Penney Jr., J.B., Young, A.B. and Newman, S.W., The globus

pallidus receives a projection from the parafascicular nucleus in the rat, 18
Kish, S.J., see Young, L.T., 323
Kishimoto, T., Sasa, M. and Takaori, S., Inhibition of lateral vestibular nucleus neurons by 5-hydroxytryptamine derived from the dorsal raphe nucleus, 229
Kogure, K., see Kato, H., 238
Kogure, K., see Kato, H., 33
Kozlowski, M.R., Sandler, P., Lin, P.-F. and Watson, A., Brain-derived cells contain a specific binding site for Gp120 which is not the CD4 antigen, 300
Kuhlmann, D., see Madeja, M., 27

Lachman, H.M., see Weiner, E.D., 117
Lacombe, P., see Dauphin, F., 75
Laguzzi, R., see Orer, H.S., 123
Lahtinen, H., see Ylinen, A., 195
Launay, J.-M., see Chapouthier, G., 342
Lee, K.S., Brooks, P. and Lowenkopf, T., Transient ischemia attenuates neuronal afterdischarges induced in the absence of synaptic transmission, 171
Lees, J., see Hynes, N., 4
Leung, L.-W.S. and Yim, C.-Y.C., Intrinsic membrane potential oscillations in hippocampal neurons in vitro, 261
Levine, J.D., see Miaskowski, C., 105
Li, P.P., see Young, L.T., 323
Li, W. and Hexum, T.D., Characterization of neuropeptide Y (NPY) receptors in human hippocampus, 167
Li, Y.-P. and Lombardini, J.B., Inhibition by taurine of the phosphorylation of specific synaptosomal proteins in the rat cortex: effects of taurine on the stimulation of calcium uptake in mitochondria and inhibition of phosphoinositide turnover, 89
Lin, P.-F., see Kozlowski, M.R., 300
Liu, G., see Huang, C.-m., 201
Liu, G., see Huang, C., 129
Liu, Y., see Kato, H., 238
Lochner, A., see Amiri', Z., 155
Lombardini, J.B., see Li, Y.-P., 89
Löscher, W., see Wahnschaffe, U., 181
Louilot, A., Gonzalez-Mora, J.L., Guadalupe, T. and Mas, M., Sex-related olfactory stimuli induce a selective increase in dopamine release in the nucleus accumbens of male rats. A voltammetric study, 313
Lowenkopf, T., see Lee, K.S., 171

Maciewicz, R., see Davar, G., 327
Madeja, M., Mußhoff, U., Kuhlmann, D. and Speckmann, E.-J., Membrane currents elicited by the epileptogenic drug pentylenetetrazole in the native oocyte of *Xenopus laevis*, 27
Martin, B., see Chapouthier, G., 342
Mas, M., see Louilot, A., 313
Matsuoka, N., Kaneko, S. and Satoh, M., Somatostatin augments long-term potentiation of the mossy fiber-CA3 system in guinea-pig hippocampal slices, 188
McCulloch, J., see Dewar, D., 58
Melograna, J.M., see Gibson, R.E., 110
Merahi, N., see Orer, H.S., 123
Miaskowski, C., Sutters, K.A., Taiwo, Y.O. and Levine, J.D., Comparison of the antinociceptive and motor effects of intrathecal opioid agonists in the rat, 105
Minciacchi, D., Granato, A. and Barbaresi, P., Organization of claustro-cortical projections to the primary somatosensory area of primates, 309
Miyake, T., see Sasaki, S.-I., 159
Morari, M., Calo, G., Antonelli, T., Gaist, G., Acciarri, N., Fabrizi, A., Bianchi, C. and Beani, L., Inhibitory effect of NMDA receptor activation on quisqualate-stimulated phosphatidylinositol turnover in the human cerebral cortex, 14
Mußhoff, U., see Madeja, M., 27

Nakano, Y., Suda, T., Sumitomo, T., Tozawa, F. and Demura, H., Effects of sex steroids on β-endorphin release from rat hypothalamus in vitro, 1
Nakazawa, K., Inoue, K., Ohara-Imaizumi, M., Fujimori, K. and Takanaka, A., Inhibition of Ca-channels by diazepam compared with that by nicardipine in pheochromocytoma PC12 cells, 44
Newman, S.W., see Kincaid, A.E., 18
Nosjean, A., see Orer, H.S., 123

Ohara-Imaizumi, M., see Nakazawa, K., 44
Orer, H.S., Merahi, N., Nosjean, A., Fattaccini, C.-M. and Laguzzi, R., Cardiovascular effects of the local injection of 5,7-dihydroxytryptamine into the nodose ganglia and nucleus tractus solitarius in awake freely moving rats, 123
Ormandy, G.C. and Jope, R.S., Pertussis toxin potentiates seizures induced by pilocarpine, kainic acid and *N*-methyl-D-aspartate, 51
Osborne, N.N., Inhibition of cAMP production by α_2-adrenoceptor stimulation in rabbit retina, 84

Papolos, D.F., see Weiner, E.D., 117
Partanen, J., see Ylinen, A., 195
Penney Jr., J.B., see Kincaid, A.E., 18
Peroutka, S.J., Cortical and striatal variations in drug competition studies with putative 5-hydroxytryptamine$_{1D}$ binding sites, 206
Phillis, J.W., see Simpson, R.E., 305
Pincus, D.W., Diciccobloom, E. and Black, I.B., Role of voltage-sensitive calcium channels in mitogenic stimulation of neuroblasts, 211
Powley, T.L., see Berthoud, H.-R., 336
Puig, S., Rivot, J.P. and Besson, J.M., In vivo electrochemical evidence that the tricyclic antidepressant femoxetine potentiates the morphine-induced increase in 5-HT metabolism in the medullary dorsal horn of freely moving rats, 222

Randić, M., see Kangrga, I., 347
Reba, R.C., see Gibson, R.E., 110
Riekkinen, P., see Ylinen, A., 195
Rivot, J.P., see Puig, S., 222
Robertson, D.W., see Gehlert, D.R., 149
Rothwell, N.J., see Strijbos, P.J.L.M., 249
Ruch, J., see Gibson, R.E., 110

Sandler, P., see Kozlowski, M.R., 300
Sano, Y., see Ihara, N., 39
Sasa, M., see Kishimoto, T., 229
Sasaki, S.-I., Uchino, H., Imagawa, M., Miyake, T. and Uchino, Y., Lower lumbar branching of caudal medullary expiratory neurons of the cat, 159
Satoh, M., see Matsuoka, N., 188
Schul, R. and Frenk, H., The role of serotonin in analgesia elicited by morphine in the periaqueductal gray matter (PAG), 353
Schwarting, R.K.W., see Adams, F., 318
Sercombe, R., see Dauphin, F., 75
Seylaz, J., see Dauphin, F., 75
Shults, C.W., see Carman, L.S., 275
Simpson, R.E. and Phillis, J.W., Adenosine deaminase reduces hypoxic and hypercapnic dilatation of rat pial arterioles: evidence for mediation by adenosine, 305
Sirviö, J., see Ylinen, A., 195
Siu, K.P., see Young, L.T., 323
Smith, G.P., see Corp, E.S., 175
Smith, S.L., see Hall, E.D., 243
Speckmann, E.-J., see Madeja, M., 27
Strijbos, P.J.L.M., Tilders, F.J.H., Carey, F., Forder, R. and Rothwell, N.J., Localization of immunoreactive lipocortin-1 in the brain and pituitary gland of the rat. Effects of adrenalectomy, dexamethasone and colchicine treatment, 249
Suda, T., see Nakano, Y., 1
Sumitomo, T., see Nakano, Y., 1
Sutters, K.A., see Miaskowski, C., 105

Taiwo, Y.O., see Miaskowski, C., 105
Takanaka, A., see Nakazawa, K., 44
Takaori, S., see Kishimoto, T., 229
Tanabe, T., see Ihara, N., 39
Tejwani, G.A., see Gudehithlu, K.P., 284
Tilders, F.J.H., see Strijbos, P.J.L.M., 249
Tozawa, F., see Nakano, Y., 1
Treiman, D.M., see Walton, N.Y., 97

Uchino, Y., see Sasaki, S.-I., 159
Uchino, H., see Sasaki, S.-I., 159
Ueda, S., see Ihara, N., 39

Vos, B., see Davar, G., 327

Wahnschaffe, U. and Löscher, W., Anticonvulsant effects of ipsilateral but not contralateral microinjections of the dopamine D_2 agonist LY 171555 into the nucleus accumbens of amygdala-kindled rats, 181
Walicke, P.A., see Frautschy, S.A., 291
Walton, N.Y. and Treiman, D.M., Motor and electroencephalographic response of refractory experimental status epilepticus in rats to treatment with MK-801, diazepam, or MK-801 plus diazepam, 97
Wang, T.F., see Gibson, R.E., 110
Warsh, J.J., see Young, L.T., 323
Watson, A., see Kozlowski, M.R., 300
Weiner, E.D., Kalasapudi, V.D., Papolos, D.F. and Lachman, H.M., Lithium aug-

ments pilocarpine-induced *fos* gene expression in rat brain, 117
Weizman, R., see Amiri', Z., 155
Wessendorf, M.W., Fluoro-Gold: composition, and mechanism of uptake, 135
Widzowski, D.V., see Cory-Slechta, D.A., 65
Wong, D.T., see Gehlert, D.R., 149
Yim, C.-Y.C., see Leung, L.-W.S., 261
Ylinen, A., Lahtinen, H., Sirviö, J., Partanen, J., Asikainen, A., Gulyas, A., Freund, T.F. and Riekkinen, P., Behavioural, electrophysiological and histopathological changes following sustained stimulation of the perforant pathway input to the hippocampus: effect of the NMDA receptor antagonist, CGP 39551, 195
Young, A.B., see Kincaid, A.E., 18
Young, L.T., Li, P.P., Kish, S.J., Siu, K.P. and Warsh, J.J., Postmortem cerebral cortex G_s α-subunit levels are elevated in bipolar affective disorder, 323
Zeeberg, B.R., see Gibson, R.E., 110

(contents continued)

Cell migration along glial fibers in dissociated cell culture of the frog optic tectum
T. Becker and C.G. Becker (F.R.G.) 331
Morphology and distribution of efferent vagal innervation of rat pancreas as revealed with anterograde transport of DiI
H.-R. Berthoud and T.L. Powley (U.S.A.) 336
Genetic difference in sensitivity to β-carboline: evidence for the involvement of brain benzodiazepine receptors
G. Chapouthier, D. Bondoux, B. Martin, C. Desforges and J.-M. Launay (France) 342
Outflow of endogenous aspartate and glutamate from the rat spinal dorsal horn in vitro by activation of low- and high-threshold primary afferent fibers. Modulation by μ-opioids
I. Kangrga and M. Randić (U.S.A.) 347
The role of serotonin in analgesia elicited by morphine in the periaqueductal gray matter (PAG)
R. Schul and H. Frenk (Israel) 353

Author Index 358